

EXHIBIT 2

IN THE UNITED STATES DISTRICT COURT
FOR THE SOUTHERN DISTRICT OF WEST VIRGINIA
CHARLESTON DIVISION

B.P.J., by her next friend and mother,
HEATHER JACKSON,

Plaintiff,

vs.

WEST VIRGINIA STATE BOARD OF
EDUCATION; HARRISON COUNTY BOARD
OF EDUCATION; WEST VIRGINIA
SECONDARY SCHOOLS ACTIVITIES
COMMISSION; W. CLAYTON BURCH, in his
official capacity as State Superintendent, DORA
STUTLER, in her official capacity as the
Harrison County Superintendent, and the
STATE OF WEST VIRGINIA,

Defendants,

and

LAINEY ARMISTEAD,

Defendant-Intervenor.

Case No. 2:21-cv-00316

Hon. Joseph R. Goodwin

DECLARATION OF GREGORY A. BROWN, PH.D., FACSM

I, Dr. Gregory A. Brown, pursuant to 28 U.S. Code § 1746, declare under penalty of perjury under the laws of the United States of America that the facts contained in my Expert Declaration of Gregory A. Brown, Ph.D., FACSM in the Case of B.P.J. v. West Virginia State Board of Education, attached hereto, are true and correct to the best of my knowledge and belief, and that the opinions expressed therein represent my own expert opinions.

Executed on February 23, 2022.



Gregory A. Brown

G. Brown

Expert Report, B.P.J. v. WV BOE et al.

Expert Report of

Gregory A Brown, Ph.D. FACSM

In the case of B.P.J. vs. West Virginia State Board of Education.

Overview

In this declaration, I explore three important questions relevant to current discussions and policy decisions concerning inclusion of transgender individuals in women's athletic competitions. Based on my professional familiarity with exercise physiology and my review of the currently available science, including that contained in the many academic sources I cite in this report, I set out and explain three basic conclusions:

- At the level of (a) elite, (b) collegiate, (c) scholastic, and (d) recreational competition, men, adolescent boys, or male children, have an advantage over equally aged, gifted, and trained women, adolescent girls, or female children in almost all athletic events;
- Biological male physiology is the basis for the performance advantage that men, adolescent boys, or male children have over women, adolescent girls, or female children in almost all athletic events; and
- The administration of androgen inhibitors and cross-sex hormones to men or adolescent boys after the onset of male puberty does not eliminate the performance advantage that men and adolescent boys have over women and adolescent girls in almost all athletic events. Likewise, there is no published scientific evidence that the administration of puberty blockers to males before puberty eliminates the pre-existing athletic advantage that prepubertal males have over prepubertal females in almost all athletic events.

In short summary, men, adolescent boys, and prepubertal male children perform better in almost all sports than women, adolescent girls, and prepubertal female children because of their inherent physiological advantages. In general, men, adolescent boys, and prepubertal male children, can run faster, output more muscular power, jump higher, and possess greater muscular endurance than women, adolescent girls, and prepubertal female children. These advantages become greater during and after male puberty, but they exist before puberty.

Further, while after the onset of puberty males are on average taller and heavier than females, a male performance advantage over females has been measured in weightlifting competitions even between males and females matched for body mass.

Male advantages in measurements of body composition, tests of physical fitness, and athletic performance have also been shown in children before puberty. These advantages are magnified during puberty, triggered in large part by the higher testosterone concentrations in men, and adolescent boys, after the onset of

adolescent girls and prepubertal female children, in competitions involving running speed, swimming speed, cycling speed, jumping height, jumping distance, and strength (to name a few, but not all, of the performance differences). As I discuss later, it is now clear that these performance advantages for men, adolescent boys, and prepubertal male children, are inherent to the biological differences between the sexes.

10. In fact, I am not aware of any scientific evidence today that disproves that after puberty men possess large advantages in athletic performance over women—so large that they are generally insurmountable for comparably gifted and trained athletes at every level (i.e. (a) elite, (b) collegiate, (c) scholastic, and (d) recreational competition). And I am not aware of any scientific evidence today that disproves that these measured performance advantages are at least largely the result of physiological differences between men and women which have been measured and are reasonably well understood.

11. My use of the term “advantage” in this paper must not be read to imply any normative judgment. The adult female physique is simply different from the adult male physique. Obviously, it is optimized in important respects for the difficult task of childbearing. On average, women require far fewer calories for healthy survival. Evolutionary biologists can and do theorize about the survival value or “advantages” provided by these and other distinctive characteristics of the female physique, but I will leave that to the evolutionary biologists. I use “advantage” to refer merely to performance advantages in athletic competitions.

12. I find in the literature a widespread consensus that the large performance and physiological advantages possessed by males—rather than social considerations or considerations of identity—are precisely the *reason* that most athletic competitions are separated by sex, with women treated as a “protected class.” To cite only a few statements accepting this as the justification:

- Handelsman et al. (2018) wrote, “Virtually all elite sports are segregated into male and female competitions. The main justification is to allow women a chance to win, as women have major disadvantages against men who are, on average, taller, stronger, and faster and have greater endurance due to their larger, stronger, muscles and bones as well as a higher circulating hemoglobin level.” (803)
- Millard-Stafford et al. (2018) wrote “Current evidence suggests that women will not swim or run as fast as men in Olympic events, which speaks against eliminating sex segregation in these individual sports” (530) “Given the historical context (2% narrowing in swimming over 44 y), a reasonable assumption might be that no more than 2% of the

current performance gap could still potentially be attributed to sociocultural influences.”, (533) and “Performance gaps between US men and women stabilized within less than a decade after federal legislation provided equal opportunities for female participation, but only modestly closed the overall gap in Olympic swimming by 2% (5% in running).” (533) Dr. Millard-Stafford, a full professor at Georgia Tech, holds a Ph.D. in Exercise Physiology and is a past President of the American College of Sports Medicine.

- In 2021, Hilton et al. wrote, “most sports have a female category the purpose of which is the protection of both fairness and, in some sports, safety/welfare of athletes who do not benefit from the physiological changes induced by male levels of testosterone from puberty onwards.” (204)
- In 2020 the Swiss High Court (“Tribunal Fédéral”) observed that “in most sports . . . women and men compete in two separate categories, because the latter possess natural advantages in terms of physiology.”¹
- The members of the Women’s Sports Policy Working Group wrote that “If sports were not sex-segregated, female athletes would rarely be seen in finals or on victory podiums,” and that “We have separate sex sport and eligibility criteria based on biological sex because this is the only way we can assure that female athletes have the same opportunities as male athletes not only to participate but to win in competitive sport. . . . If we did not separate athletes on the basis of biological sex—if we used any other physical criteria—we would never see females in finals or on podiums.” (WSPWG Briefing Book 2021 at 5, 20.)
- In 2020, the World Rugby organization stated that “the women’s category exists to ensure protection, safety and equality for those who do not benefit from the biological advantage created by these biological performance attributes.” (World Rugby Transgender Women Guidelines 2020.)
- In 2021 Harper et al. stated “...the small decrease in strength in transwomen after 12–36 months of GAHT [Gender Affirming Hormone Therapy] suggests that transwomen likely retain a strength advantage

¹ “dans la plupart des sports . . . les femmes et les hommes concourent dans deux catégories séparées, ces derniers étant naturellement avantagés du point de vue physique.” Tribunal Fédéral decision of August 25, 2020, Case 4A_248/2019, 4A_398/2019, at §9.8.3.3.

over cisgender women.” (7) and “...observations in trained transgender individuals are consistent with the findings of the current review in untrained transgender individuals, whereby 30 months of GAHT may be sufficient to attenuate some, but not all, influencing factors associated with muscular endurance and performance.” (8)

- Hamilton et al. (2021), in a consensus statement for the International Federation of Sports Medicine (FIMS) concluded that “Transwomen have the right to compete in sports. However, cisgender women have the right to compete in a protected category.” (1409)

13. While the sources I mention above gather more extensive scientific evidence of this uncontroversial truth, I provide here a brief summary of representative facts concerning the male advantage in athletic performance.

A. Men are stronger.

14. Males exhibit greater strength throughout the body. Both Handelsman et al. (2018) and Hilton & Lundberg (2021) have gathered multiple literature references that document this fact in various muscle groups.

15. Men have in the neighborhood of 60%-100% greater **arm strength** than women. (Handelsman 2018 at 812.)² One study of elbow flexion strength (basically, bringing the fist up towards the shoulder) in a large sample of men and women found that men exhibited 109% greater isometric strength, and 89% higher strength in a single repetition. (Hilton 2021 at 204, summarizing Hubal (2005) at Table 2.)

16. **Grip strength** is often used as a useful proxy for strength more generally. In one study, men showed on average 57% greater grip strength than women. (Bohannon 2019.) A wider meta-analysis of multiple grip-strength studies not limited to athletic populations found that 18- and 19-year-old males exhibited in

² Handelsman expresses this as women having 50% to 60% of the “upper limb” strength of men. Handelsman cites Sale, *Neuromuscular function*, for this figure and the “lower limb” strength figure. Knox et al., *Transwomen in elite sport* (2018) are probably confusing the correct way to state percentages when they state that “differences lead to decreased trunk and lower body strength by 64% and 72% respectively, in women” (397): interpreted literally, this would imply that men have **almost 4x as much** lower body strength as do women.

the neighborhood of 2/3 greater grip strength than females. (Handelsman 2017 Figure 3, summarizing Silverman 2011 Table 1.)³

17. In an evaluation of maximal isometric handgrip strength in 1,654 healthy men, 533 healthy women aged 20-25 years and 60 “highly trained elite female athletes from sports known to require high hand-grip forces (judo, handball),” Leyk et al. (2007) observed that, “The results of female national elite athletes even indicate that the strength level attainable by extremely high training will rarely surpass the 50th percentile of untrained or not specifically trained men.” (Leyk 2007 at 415.)

18. Men have in the neighborhood of 25%-60% greater **leg strength** than women. (Handelsman 2018 at 812.) In another measure, men exhibit 54% greater knee extension torque and this male leg strength advantage is consistent across the lifespan. (Neder 1999 at 120-121.)

19. When male and female Olympic weightlifters of the same body weight are compared, the top males lift weights between 30% and 40% greater than the females of the same body weight. But when top male and female performances are compared in powerlifting, without imposing any artificial limitations on bodyweight, the male record is 65% higher than the female record. (Hilton 2021 at 203.)

20. In another measure that combines many muscle groups as well as weight and speed, moderately trained males generated 162% greater punching power than females even though men do not possess this large an advantage in any single bio-mechanical variable. (Morris 2020.) This objective reality was subjectively summed up by women’s mixed-martial arts fighter Tamikka Brents, who suffered significant facial injuries when she fought against a biological male who identified as female and fought under the name of Fallon Fox. Describing the experience, Brents said:

“I’ve fought a lot of women and have never felt the strength that I felt in a fight as I did that night. I can’t answer whether it’s because she was born a man or not because I’m not a doctor. I can only say, I’ve never felt so overpowered ever in my life, and I am an abnormally strong female in my own right.”⁴

³ Citing Silverman, *The secular trend for grip strength in Canada and the United States*, J. Ports Sci. 29:599-606 (2011).

⁴ <http://whoatv.com/exclusive-fallon-foxs-latest-opponent-opens-up-to-whoatv/> (last accessed October 5, 2021).


B. Men run faster.

21. Many scholars have detailed the wide performance advantages enjoyed by men in running speed. One can come at this reality from a variety of angles.

22. Multiple authors report a male speed advantage in the neighborhood of 10%-13% in a variety of events, with a variety of study populations. Handelsman et al. 2018 at 813 and Handelsman 2017 at 70 both report a male advantage of about 10% by age 17. Thibault et al. 2010 at 217 similarly reported a stable 10% performance advantage across multiple events at the Olympic level. Tønnesen et al. (2015 at 1-2) surveyed the data and found a consistent male advantage of 10%-12% in running events after the completion of puberty. They document this for both short sprints and longer distances. One group of authors found that the male advantage increased dramatically in ultra-long-distance competition (Lepers & Knechtle 2013.)

23. A great deal of current interest has been focused on track events. It is worth noting that a recent analysis of publicly available sports federation and tournament records found that men enjoy the *least* advantage in running events, as compared to a range of other events and metrics, including jumping, pole vaulting, tennis serve speed, golf drives, baseball pitching speed, and weightlifting. (Hilton 2021 at 201-202.) Nevertheless, as any serious runner will recognize, the approximately 10% male advantage in running is an overwhelming difference. Dr. Hilton calculates that “approximately 10,000 males have personal best times that are faster than the current Olympic 100m female champion.” (Hilton 2021 at 204.) Professors Doriane Coleman, Jeff Wald, Wickliffe Shreve, and Richard Clark dramatically illustrated this by compiling the data and creating the figure below (last accessed on February 10, 2022, at <https://bit.ly/35yOyS4>), which shows that the *lifetime best performances* of three female Olympic champions in the 400m event—including Team USA’s Sanya Richards-Ross and Allyson Felix—would not match the performances of “literally thousands of boys and men, including thousands who would be considered second tier in the men’s category” *just in 2017 alone*: (data were drawn from the International Association of Athletics Federations (IAAF) website which provides complete, worldwide results for individuals and events, including on an annual and an all-time basis).

Comparing the Best Elite Females to Boys and Men:
Personal Bests for 3 Female Gold Medalists versus 2017 Performances by Boys and Men



24. Professor Coleman and her colleague Wicklyffe Shreve also created the table below (last accessed on February 10, 2022, at <https://bit.ly/37E1s2X>), which “compares the number of men—males over 18—competing in events reported to the International Association of Athletics Federation whose results in each event in 2017 would have ranked them above the very best elite woman that year.”

TABLE 2 – World’s Best Woman v. Number of Men Outperforming			
Event	Best Women’s Result	Best Men’s Result	# of Men Outperforming
100 Meters	10.71	9.69	2,474
200 Meters	21.77	19.77	2,920
400 Meters	49.46	43.62	4,341
800 Meters	1:55.16*	1:43.10	3,992+
1500 Meters	3:56.14	3:28.80	3,216+
3000 Meters	8:23.14	7:28.73	1307+
5000 Meters	14:18.37	12:55.23	1,243
High Jump	2.06 meters	2.40 meters	777
Pole Vault	4.91 meters	6.00 meters	684
Long Jump	7.13 meters	8.65 meters	1,652
Triple Jump	14.96 meters	18.11 meters	969

25. The male advantage becomes insuperable well before the developmental changes of puberty are complete. Dr. Hilton documents that even “schoolboys”—defined as age 15 and under—have beaten the female world records in running, jumping, and throwing events. (Hilton 2021 at 204.)

26. Similarly, Coleman and Shreve created the table below (last accessed on February 10, 2022, at <https://bit.ly/37E1s2X>), which “compares the number of boys—males under the age of 18—whose results in each event in 2017 would rank them above the single very best elite [adult] woman that year;” data were drawn from the International Association of Athletics Federations (IAAF) website

TABLE 1 – World’s Best Woman v. Under 18 Boys			
Event	Best Women’s Result	Best Boys’ Result	# of Boys Outperforming
100 Meters	10.71	10.15	124 ⁺
200 Meters	21.77	20.51	182
400 Meters	49.46	45.38	285
800 Meters	1:55.16*	1:46.3	201+
1500 Meters	3:56.14	3:37.43	101+
3000 Meters	8:23.14	7:38.90	30
5000 Meters	14:18.37	12:55.58	15
High Jump	2.06 meters	2.25 meters	28
Pole Vault	4.91 meters	5.31 meters	10
Long Jump	7.13 meters	7.88 meters	74
Triple Jump	14.96 meters	17.30 meters	47

27. In an analysis I have performed of running events (consisting of the 100 m, 200 m, 400 m, 800 m, 1500 m, 5000 m, and 10000 m) in the Division 1, Division 2, and Division 3 NCAA Outdoor track championships for the years of 2010-2019, the average performance across all events of the 1st place man was 14.1% faster than the 1st place woman, with the smallest difference being a 10.2% advantage for men in the Division 1 100 m race. The average 8th place man across all events (the last place to earn the title of All American) was 11.2% faster than 1st place woman, with the smallest difference being a 6.5% advantage for men in the Division 1 100 m race. (Brown et al. Unpublished observations, to be presented at the 2022 Annual Meeting of the American College of Sports Medicine.)

28. Athletic.net® is an internet-based resource providing “results, team, and event management tools to help coaches and athletes thrive.” Among the resources available on Athletic.net are event records that can be searched by nationally or by state age group, school grade, and state. Higerd (2021) in an evaluation of high school track running performance records from five states(CA, FL, MN, NY, WA), over three years (2017 – 2019) observed that males were 14.38% faster than females in the 100M (at 99), 16.17% faster in the 200M (at 100), 17.62% faster in the 400M (at 102), 17.96% faster in the 800M (at 103), 17.81% faster in the 1600M (at 105), and 16.83% faster in the 3200M (at 106).

C. Men jump higher and farther.

29. Jumping involves both leg strength and speed as positive factors, with body weight of course a factor working against jump height. Despite their substantially greater body weight, males enjoy an even greater advantage in jumping than in running. Handelsman 2018 at 813, looking at youth and young adults, and Thibault 2010 at 217, looking at Olympic performances, both found male advantages in the range of 15%-20%. See also Tønnessen 2015 (approximately 19%); Handelsman 2017 (19%); Hilton 2021 at 201 (18%). Looking at the vertical jump called for in volleyball, research on elite volleyball players found that males jumped on average 50% higher during an “attack” at the net than did females. (Sattler 2015; see also Hilton 2021 at 203 (33% higher vertical jump).)

30. Higerd (2021) in an evaluation of high school high jump performance available through the track and field database athletic.net®, which included five states (CA, FL, MN, NY, WA), over three years (2017 – 2019) (at 82) observed that in 23,390 females and 26,843 males, females jumped an average of 1.35 m and males jumped an average of 1.62 m, for an 18.18% performance advantage for males (at 96). In an evaluation of long jump performance in 45,705 high school females and 54,506 high school males the females jumped an average of 4.08 m and males jumped an average of 5.20 m, for a 24.14% performance advantage for males (at 97).

31. The combined male advantage of body height and jump height means, for example, that a total of seven women in the WNBA have ever dunked a basketball in the regulation 10 foot hoop,⁵ while the ability to dunk appears to be almost universal among NBA players: “Since the 1996–97 season (the earliest data is available from Basketball-Reference.com), 1,801 different [NBA] players have combined for 210,842 regular-season dunks, and 1,259 out of 1,367 players (or 92%) who have played at least 1,000 minutes have dunked at least once.”⁶

D. Men throw, hit, and kick faster and farther.

32. Strength, arm-length, and speed combine to give men a large advantage over women in throwing. This has been measured in a number of studies.

33. One study of elite male and female baseball pitchers showed that men throw baseballs 35% faster than women—81 miles/hour for men vs. 60 miles/hour for women. (Chu 2009.) By age 12, “boys’ throwing velocity is already between 3.5 and 4 standard deviation units higher than the girls’.” (Thomas 1985 at 276.) By age seventeen, the *average* male can throw a ball farther than 99% of seventeen-year-old females. (Lombardo 2018; Chu 2009; Thomas 1985 at 268.) Looking at publicly available data, Hilton & Lundberg found that in both baseball pitching and the field hockey “drag flick,” the *record* ball speeds achieved by males are more than 50% higher than those achieved by females. (Hilton 2021 at 202-203.)

34. Men achieve serve speeds in tennis more than 15% faster than women; and likewise in golf achieve ball speeds off the tee more than 15% faster than women. (Hilton 2021 at 202.)

35. Males are able to throw a javelin more than 30% farther than females. (Lombardo 2018 Table 2; Hilton 2021 at 203.)

36. Men serve and spike volleyballs with higher velocity than women, with a performance advantage in the range of 29-34%. (Hilton 2021 at 204 Fig. 1.)

37. Men are also able to kick balls harder and faster. A study comparing collegiate soccer players found that males kick the ball with an average 20% greater velocity than females. (Sakamoto 2014.)

⁵ https://www.espn.com/wnba/story/_/id/32258450/2021-wnba-playoffs-brittney-griner-owns-wnba-dunking-record-coming-more.

⁶ <https://www.si.com/nba/2021/02/22/nba-non-dunkers-patty-mills-tj-mcconnell-steve-novak-daily-cover>

E. Males exhibit faster reaction times.

38. Interestingly, men enjoy an additional advantage over women in reaction time—an attribute not obviously related to strength or metabolism (e.g. $V_{O_2\text{max}}$). “Reaction time in sports is crucial in both simple situations such as the gun shot in sprinting and complex situations when a choice is required. In many team sports this is the foundation for tactical advantages which may eventually determine the outcome of a game.” (Dogan 2009 at 92.) “Reaction times can be an important determinant of success in the 100m sprint, where medals are often decided by hundredths or even thousandths of a second.” (Tønnessen 2013 at 885.)

39. The existence of a sex-linked difference in reaction times is consistent over a wide range of ages and athletic abilities. (Dykiert 2012.) Even by the age of 4 or 5, in a ruler-drop test, males have been shown to exhibit 4% to 6% faster reaction times than females. (Latorre-Roman 2018.) In high school athletes taking a common baseline “ImPACT” test, males showed 3% faster reaction times than females. (Mormile 2018.) Researchers have found a 6% male advantage in reaction times of both first-year medical students (Jain 2015) and world-class sprinters (Tønnessen 2013).

40. Most studies of reaction times use computerized tests which ask participants to hit a button on a keyboard or to say something in response to a stimulus. One study on NCAA athletes measured “reaction time” by a criterion perhaps more closely related to athletic performance—that is, how fast athletes covered 3.3 meters after a starting signal. Males covered the 3.3 meters 10% faster than females in response to a visual stimulus, and 16% faster than females in response to an auditory stimulus. (Spierer 2010.)

41. Researchers have speculated that sex-linked differences in brain structure, as well as estrogen receptors in the brain, may be the source of the observed male advantage in reaction times, but at present this remains a matter of speculation and hypothesis. (Mormile at 19; Spierer at 962.)

III. Men have large measured physiological differences compared to women which demonstrably or likely explain their performance advantages.

42. No single physiological characteristic alone accounts for all or any one of the measured advantages that men enjoy in athletic performance. However, scientists have identified and measured a number of physiological factors that contribute to superior male performance.

A. Men are taller and heavier than women

43. In some sports, such as basketball and volleyball, height itself provides competitive advantage. While some women are taller than some men, based on data from 20 countries in North America, Europe, East Asia, and Australia, the 50th percentile for body height for women is 164.7 cm (5 ft 5 inches) and the 50th percentile for body height for men is 178.4 cm (5 ft 10 inches). Helping to illustrate the inherent height difference between men and women, from the same data analysis, the 95th percentile for body height for women is 178.9 cm (5 feet 10.43 inches), which is only 0.5 cm taller than the 50th percentile for men (178.4 cm; 5 feet 10.24 inches), while the 95th percentile for body height for men is 193.6 cm (6 feet 4.22 inches). (Roser 2013.)

44. To look at a specific athletic population, an evaluation of NCAA Division 1 basketball players compared 68 male guards and 59 male forwards to 105 female guards and 91 female forwards, and found that on average the male guards were 187.4 ± 7.0 cm tall and weighed 85.2 ± 7.4 kg while the female guards were 171.6 ± 5.0 cm tall and weighed 68.0 ± 7.4 kg. The male forwards were 201.7 ± 4.0 cm tall and weighed 105.3 ± 5.9 kg while the female forwards were 183.5 ± 4.4 cm tall and weighed 82.2 ± 12.5 kg. (Fields 2018 at 3.)

B. Males have larger and longer bones, stronger bones, and different bone configuration.

45. Obviously, males on average have longer bones. “Sex differences in height have been the most thoroughly investigated measure of bone size, as adult height is a stable, easily quantified measure in large population samples. Extensive twin studies show that adult height is highly heritable with predominantly additive genetic effects that diverge in a sex-specific manner from the age of puberty onwards.” (Handelsman 2018 at 818.) “Pubertal testosterone exposure leads to an ultimate average greater height in men of 12–15 centimeters, larger bones, greater muscle mass, increased strength and higher hemoglobin levels.” (Gooren 2011 at 653.)

46. “Men have distinctively greater bone size, strength, and density than do women of the same age. As with muscle, sex differences in bone are absent prior to puberty but then accrue progressively from the onset of male puberty due to the sex difference in exposure to adult male circulating testosterone concentrations.” (Handelsman 2018 at 818.)

47. “[O]n average men are 7% to 8% taller with longer, denser, and stronger bones, whereas women have shorter humerus and femur cross-sectional

areas being 65% to 75% and 85%, respectively, those of men.” (Handelsman 2018 at 818.)

48. Greater height, leg, and arm length themselves provide obvious advantages in several sports. But male bone geometry also provides less obvious advantages. “The major effects of men’s larger and stronger bones would be manifest via their taller stature as well as the larger fulcrum with greater leverage for muscular limb power exerted in jumping, throwing, or other explosive power activities.” (Handelsman 2018 at 818.)

49. Male advantage in bone size is not limited to length, as larger bones provide the mechanical framework for larger muscle mass. “From puberty onwards, men have, on average, 10% more bone providing more surface area. The larger surface area of bone accommodates more skeletal muscle so, for example, men have broader shoulders allowing more muscle to build. This translates into 44% less upper body strength for women, providing men an advantage for sports like boxing, weightlifting and skiing. In similar fashion, muscle mass differences lead to decreased trunk and lower body strength by 64% and 72%, respectively in women. These differences in body strength can have a significant impact on athletic performance, and largely underwrite the significant differences in world record times and distances set by men and women.” (Knox 2019 at 397.)

50. Meanwhile, distinctive aspects of the female pelvis geometry cut against athletic performance. “[T]he widening of the female pelvis during puberty, balancing the evolutionary demands of obstetrics and locomotion, retards the improvement in female physical performance.” (Handelsman 2018 at 818.) “[T]he major female hormones, oestrogens, can have effects that disadvantage female athletic performance. For example, women have a wider pelvis changing the hip structure significantly between the sexes. Pelvis shape is established during puberty and is driven by oestrogen. The different angles resulting from the female pelvis leads to decreased joint rotation and muscle recruitment ultimately making them slower.” (Knox 2019 at 397.)

51. There are even sex-based differences in foot size and shape. Wunderlich & Cavanaugh (2001) observed that a “foot length of 257 mm represents a value that is ... approximately the 20th percentile men’s foot lengths and the 80th percentile women’s foot lengths.” (607) and “For a man and a woman, both with statures of 170 cm (5 feet 7 inches), the man would have a foot that was approximately 5 mm longer and 2 mm wider than the woman.” (608). Based on these, and other analyses, they conclude that “female feet and legs are not simply scaled-down versions of male feet but rather differ in a number of shape characteristics, particularly at the arch, the lateral side of the foot, the first toe, and the ball of the foot.” (605) Further, Fessler et al. (2005) observed that “female foot length is consistently smaller than male foot length” (44) and concludes that

“proportionate foot length is smaller in women” (51) with an overall conclusion that “Our analyses of genetically disparate populations reveal a clear pattern of sexual dimorphism, with women consistently having smaller feet proportionate to stature than men.” (53)

52. Beyond simple performance, the greater density and strength of male bones provide higher protection against stresses associated with extreme physical effort: “[S]tress fractures in athletes, mostly involving the legs, are more frequent in females, with the male protection attributable to their larger and thicker bones.” (Handelsman 2018 at 818.)

C. Males have much larger muscle mass.

53. The fact that, on average, men have substantially larger muscles than women is as well known to common observation as men’s greater height. But the male advantage in muscle size has also been extensively measured. The differential is large.

54. “On average, women have 50% to 60% of men’s upper arm muscle cross-sectional area and 65% to 70% of men’s thigh muscle cross-sectional area, and women have 50% to 60% of men’s upper limb strength and 60% to 80% of men’s leg strength. Young men have on average a skeletal muscle mass of >12 kg greater than age-matched women at any given body weight.” (Handelsman 2018 at 812. See also Gooren 2011 at 653, Thibault 2010 at 214.)

55. “There is convincing evidence that the sex differences in muscle mass and strength are sufficient to account for the increased strength and aerobic performance of men compared with women and is in keeping with the differences in world records between the sexes.” (Handelsman 2018 at 816.)

56. Once again, looking at specific and comparable populations of athletes, an evaluation of NCAA Division 1 basketball players consisting of 68 male guards and 59 male forwards, compared to 105 female guards and 91 female forwards, reported that on average the male guards had 77.7 ± 6.4 kg of fat free mass and 7.4 ± 3.1 kg fat mass while the female guards had 54.6 ± 4.4 kg fat free mass and 13.4 ± 5.4 kg fat mass. The male forwards had 89.5 ± 5.9 kg fat free mass and 15.9 ± 5.6 kg fat mass while the female forwards had 61.8 ± 5.9 kg fat free mass and 20.5 ± 7.7 kg fat mass. (Fields 2018 at 3.)

D. Females have a larger proportion of body fat.

57. While women have smaller muscles, they have proportionately more body fat, in general a negative for athletic performance. “Oestrogens also affect body

composition by influencing fat deposition. Women, on average, have higher percentage body fat, and this holds true even for highly trained healthy athletes (men 5%–10%, women 8%–15%). Fat is needed in women for normal reproduction and fertility, but it is not performance-enhancing. This means men with higher muscle mass and less body fat will normally be stronger kilogram for kilogram than women.” (Knox 2019 at 397.)

58. “[E]lite females have more (<13 vs. <5 %) body fat than males. Indeed, much of the difference in [maximal oxygen uptake] between males and females disappears when it is expressed relative to lean body mass. . . . Males possess on average 7–9 % less percent body fat than females.” (Lepers 2013 at 853.)

59. Knox et al. observe that both female pelvis shape and female body fat levels “disadvantage female athletes in sports in which speed, strength and recovery are important,” (Knox 2019 at 397), while Tønnessen et al. describe the “ratio between muscular power and total body mass” as “critical” for athletic performance. (Tønnessen 2015 at 7.)

E. Males are able to metabolize and release energy to muscles at a higher rate due to larger heart and lung size, and higher hemoglobin concentrations.

60. While advantages in bone size, muscle size, and body fat are easily perceived and understood by laymen, scientists also measure and explain the male athletic advantage at a more abstract level through measurements of metabolism, or the ability to deliver energy to muscles throughout the body.

61. Energy release at the muscles depends centrally on the body’s ability to deliver oxygen to the muscles, where it is essential to the complex chain of biochemical reactions that make energy available to power muscle fibers. Men have multiple distinctive physiological attributes that together give them a large advantage in oxygen delivery.

62. Oxygen is taken into the blood in the lungs. Men have greater capability to take in oxygen for multiple reasons. “[L]ung capacity [is] larger in men because of a lower diaphragm placement due to Y-chromosome genetic determinants.” (Knox 2019 at 397.) Supporting larger lung capacity, men have “greater cross-sectional area of the trachea”; that is, they can simply move more air in and out of their lungs in a given time. (Hilton 2021 at 201.)

63. More, male lungs provide superior oxygen exchange even for a given volume: “The greater lung volume is complemented by testosterone-driven **enhanced alveolar multiplication** rate during the early years of life. Oxygen exchange takes place between the air we breathe and the bloodstream at the alveoli,

so more alveoli allows more oxygen to pass into the bloodstream. Therefore, the greater lung capacity allows more air to be inhaled with each breath. This is coupled with an improved uptake system allowing men to absorb more oxygen.” (Knox 2019 at 397.)

64. “Once in the blood, oxygen is carried by haemoglobin. **Haemoglobin concentrations** are directly modulated by testosterone so men have higher levels and can carry more oxygen than women.” (Knox 2019 at 397.) “It is well known that levels of circulating hemoglobin are androgen-dependent and consequently higher in men than in women by 12% on average.... Increasing the amount of hemoglobin in the blood has the biological effect of increasing oxygen transport from lungs to tissues, where the increased availability of oxygen enhances aerobic energy expenditure.” (Handelsman 2018 at 816.) (See also Lepers 2013 at 853; Handelsman 2017 at 71.) “It may be estimated that as a result the average maximal oxygen transfer will be ~10% greater in men than in women, which has a direct impact on their respective athletic capacities.” (Handelsman 2018 at 816.)

65. But the male metabolic advantage is further multiplied by the fact that men are also able to **circulate more blood per second** than are women. “Oxygenated blood is pumped to the active skeletal muscle by the heart. The left ventricle chamber of the heart is the reservoir from which blood is pumped to the body. The larger the left ventricle, the more blood it can hold, and therefore, the more blood can be pumped to the body with each heartbeat, a physiological parameter called ‘stroke volume’. The female heart size is, on average, 85% that of a male resulting in the stroke volume of women being around 33% less.” (Knox 2018 at 397.) Hilton cites different studies that make the same finding, reporting that men on average can pump 30% more blood through their circulatory system per minute (“cardiac output”) than can women. (Hilton 2021 at 202.)

66. Finally, at the cell where the energy release is needed, men appear to have yet another advantage. “Additionally, there is experimental evidence that testosterone increases . . . **mitochondrial biogenesis**, myoglobin expression, and IGF-1 content, which may augment energetic and power generation of skeletal muscular activity.” (Handelsman 2018 at 811.)

67. “Putting all of this together, men have a much more efficient cardiovascular and respiratory system.” (Knox 2019 at 397.) A widely accepted measurement that reflects the combined effects of all these respiratory, cardiovascular, and metabolic advantages is referred to as “V_O2max,” which refers to the maximum rate at which an individual can consume oxygen during aerobic

outperforming girls by 100% – 1200%, highlighting the advantages in upper body strength in males.

81. A very recent literature review commissioned by the five United Kingdom governmental Sport Councils concluded that while “[i]t is often assumed that children have similar physical capacity regardless of their sex, . . . large-scale data reports on children from the age of six show that young males have significant advantage in cardiovascular endurance, muscular strength, muscular endurance, speed/agility and power tests,” although they “score lower on flexibility tests.” (UK Sports Councils’ Literature Review 2021 at 3.)

82. Hilton et al., also writing in 2021, reached the same conclusion: “An extensive review of fitness data from over 85,000 Australian children aged 9–17 years old showed that, compared with 9-year-old females, 9-year-old males were faster over short sprints (9.8%) and 1 mile (16.6%), could jump 9.5% further from a standing start (a test of explosive power), could complete 33% more push-ups in 30 [seconds] and had 13.8% stronger grip.” (Hilton 2021 at 201, summarizing the findings of Catley & Tomkinson 2013.)

83. The following data are taken from Catley & Tomkinson (2013 at 101) showing the low, middle, and top decile for 1.6 km run (1.0 mile) run time for 11,423 girls and boys ages 9-17.

1.6 km run (1.0 mile) run time for 11,423 girls and boys ages 9-17

Age	Male			Female			Male-Female % Difference		
	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile
9	684	522	423	769.0	609.0	499.0	11.1%	14.3%	15.2%
10	666	511	420	759.0	600.0	494.0	12.3%	14.8%	15.0%
11	646	500	416	741.0	586.0	483.0	12.8%	14.7%	13.9%
12	621	485	408	726.0	575.0	474.0	14.5%	15.7%	13.9%
13	587	465	395	716.0	569.0	469.0	18.0%	18.3%	15.8%
14	556	446	382	711.0	567.0	468.0	21.8%	21.3%	18.4%
15	531	432	373	710.0	570.0	469.0	25.2%	24.2%	20.5%
16	514	423	366	710.0	573.0	471.0	27.6%	26.2%	22.3%
17	500	417	362	708.0	575.0	471.0	29.4%	27.5%	23.1%

84. Tomkinson et al. (2018) performed a similarly extensive analysis of literally millions of measurements of a variety of strength and agility metrics from the “Eurofit” test battery on children from 30 European countries. They provide detailed results for each metric, broken out by decile. Sampling the low, middle, and top decile, 9-year-old boys performed better than 9-year-old girls by between 6.5%

and 9.7% in the standing broad jump; from 11.4% to 16.1% better in handgrip; and from 45.5% to 49.7% better in the “bent-arm hang.” (Tomkinson 2018.)

85. The Bent Arm Hang test is a measure of upper body muscular strength and endurance used in the Eurofit Physical Fitness Test Battery. To perform the Bent Arm Hang, the child is assisted into position with the body lifted to a height so that the chin is level with the horizontal bar (like a pull up bar). The bar is grasped with the palms facing away from body and the hands shoulder width apart. The timing starts when the child is released. The child then attempts to hold this position for as long as possible. Timing stops when the child's chin falls below the level of the bar, or the head is tilted backward to enable the chin to stay level with the bar.

86. Using data from Tomkinson (2018; table 7 at 1452), the following table sampling the low, middle, and top decile for bent arm hang for 9- to 17-year-old children can be constructed:

Bent Arm Hang time (in seconds) for children ages 9 - 17 years

Age	Male			Female			Male-Female % Difference		
	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile	10th %ile	50th %ile	90th %ile
9	2.13	7.48	25.36	1.43	5.14	16.94	48.95%	45.53%	49.70%
10	2.25	7.92	26.62	1.42	5.15	17.06	58.45%	53.79%	56.04%
11	2.35	8.32	27.73	1.42	5.16	17.18	65.49%	61.24%	61.41%
12	2.48	8.79	28.99	1.41	5.17	17.22	75.89%	70.02%	68.35%
13	2.77	9.81	31.57	1.41	5.18	17.33	96.45%	89.38%	82.17%
14	3.67	12.70	38.39	1.40	5.23	17.83	162.14%	142.83%	115.31%
15	5.40	17.43	47.44	1.38	5.35	18.80	291.30%	225.79%	152.34%
16	7.39	21.75	53.13	1.38	5.63	20.57	435.51%	286.32%	158.29%
17	9.03	24.46	54.66	1.43	6.16	23.61	531.47%	297.08%	131.51%

87. Evaluating these data, a 9-year-old boy in the 50th percentile (that is to say a 9-year-old boy of average upper body muscular strength and endurance) will perform better in the bent arm hang test than 9 through 17-year-old girls in the 50th percentile. Similarly, a 9-year-old boy in the 90th percentile will perform better in the bent arm hang test than 9 through 17-year-old girls in the 90th percentile.

88. Using data from Tomkinson et al. (2017; table 1 at 1549), the following table sampling the low, middle, and top decile for running speed in the last stage of the 20 m shuttle run for 9- to 17-year-old children can be constructed.